



Clonidine

0.1mg/mL Oral Suspension

Last updated: April 2007

Batch No: _____

Disclaimers: CAUTION: The information on The Hospital for Sick Children World Wide Web Site has been developed specifically for The Hospital for Sick Children. It is provided for informational purposes for qualified health care professionals. This material is not intended as a substitute for consulting qualified health care professionals. Patient circumstances will vary and some information may have become outdated as a result of more recent medical developments.

Ingredients	Mfr	Lot #	Expiry Date	Quantity	Measured	Checked
Clonidine 0.2 mg Tablets*	BOE			30		
Sterile Water for Irrigation USP				2mL		
Simple Syrup NF/BP/USP				qs 60mL		

Equipment

- glass mortar and pestle
- 100mL graduate
- glass stirring rod
- 3mL oral syringe
- 100mL amber bottle
- shrink seal

✓ Procedure

1. Place Clonidine tablets in a glass mortar
2. Soak tablets with sterile water and small amount of Simple Syrup for about 15-20 minutes. Then crush tablets.
3. Add more Simple Syrup in small amounts to get a smooth paste. Mix well after each addition until a liquid is formed.
4. Transfer to graduate. Rinse mortar out 2-3 times with Simple Syrup to ensure all drug is transferred to graduate
5. Mix well. Transfer to clean, glass amber bottle
6. Label. Affix seals if necessary

Storage: Refrigerate

Expiry: 28 days

Sample Label:

	CLONIDINE 0.1MG/ML ORAL SUSPNSION
L#:	EXP:
REFRIGERATE	SHAKE WELL (60ML)

Date and Signature (Technician and Pharmacist Final Check): _____
