



Flecainide Acetate

20mg/mL Oral Suspension

Last updated: April 2007

Batch No: _____

Disclaimers: CAUTION: The information on The Hospital for Sick Children World Wide Web Site has been developed specifically for The Hospital for Sick Children. It is provided for informational purposes for qualified health care professionals. This material is not intended as a substitute for consulting qualified health care professionals. Patient circumstances will vary and some information may have become outdated as a result of more recent medical developments.

| Ingredients | Mfr | Lot # | Expiry Date | Quantity | Measured | Checked |
|----------------------------------|----------|-------|-------------|----------|----------|---------|
| Flecainide acetate 100mg tablets | 3M | | | 12 | | |
| Ora-Plus® | Paddock | | | 30mL | | |
| *Ora-Sweet® | Paddock | | | 30mL | | |
| Ora-Plus®/Ora-Sweet® Mixture | SickKids | | | qs 60mL | | |

*Can substitute with Ora-Sweet SF® for a sugar-free suspension.

Equipment

- Glass mortar and pestle
- 125mL graduated measure x2
- Glass stirring rod
- 100mL clean amber bottle
- Shrink seal

✓ Procedure

1. Measure out and mix together the Ora-Plus® and Ora-Sweet® vehicles.
2. Put the flecainide acetate tablets into the mortar and crush into a fine powder.
3. Add Ora-Plus®/Ora-Sweet® mixture in small amounts to the mortar until a smooth paste is formed. Add more vehicle to the paste until a liquid is formed. Transfer contents into the graduate.
4. Use additional vehicle to rinse the remaining drug from the mortar and add to the graduate.
5. Make up to the final volume of 60mL using vehicle. Stir well.
6. Transfer the suspension into the amber bottle.
7. Label and affix shrink seal.

Storage: Room temperature

Expiry: 60 days

Sample Label:

| | |
|------------------|--|
| | FLECAINIDE ACETATE 20MG/ML SUSPENSION |
| L#: | EXP: |
| ROOM TEMPERATURE | SHAKE WELL (60ML) |

Date and Signature (Technician and Pharmacist Final Check): _____
