



SickKids Mouthwash For Pain

Last updated: April 2007

Batch No: _____

Disclaimers: CAUTION: The information on The Hospital for Sick Children World Wide Web Site has been developed specifically for The Hospital for Sick Children. It is provided for informational purposes for qualified health care professionals. This material is not intended as a substitute for consulting qualified health care professionals. Patient circumstances will vary and some information may have become outdated as a result of more recent medical developments.

Ingredients	Mfr	Lot #	Expiry Date	Quantity	Measured	Checked
Sugar-Free Kool-Aid Crystals (Tropical Punch Flavour)				4.5 gm (1pkg)		
Sterile Water for Irrigation USP				qs1500mL		
Lidocaine HCl Viscous 2%				1000mL		
Sugar-Free Kool-Aid Solution (see steps 1 & 2)				qs1700mL		

Equipment

- 3000mL plastic graduated measure
- 5000mL plastic graduated measure
- glass stirring rod
- 25mL clean amber bottles
- 100mL clean amber bottles
- shrink seals

✓ Procedure

1. Empty the contents of the Kool-Aid powder packet into the 3000mL graduated measure.
2. Add approximately 500mL of Sterile Water for Irrigation to the powder and stir well until dissolved.
3. Once dissolved make up to the final volume of 1500mL with Sterile Water for Irrigation. (set aside)
4. Measure 1000mL of Lidocaine Viscous 2% into the 3000mL graduated measure.
5. Add enough Kool-Aid solution to the Lidocaine Viscous to make up to the volume of 1700mL. Stir well until mixed.
6. Pour mouthwash into the appropriate size bottles
7. Label and affix shrink seals.

Storage: Room temperature

Expiry: 2 months

Sample Label:

	MOUTHWASH FOR PAIN
	EACH 1ML CONTAINS : LIDOCAINE HCL VISCOUS 12MG IN SUGAR FREE KOOLAID
L#:	EXP:
ROOM TEMPERATURE	SHAKE WELL (100ML)

Date and Signature (Technician and Pharmacist Final Check): _____