



Sotalol HCL

5mg/mL Oral Suspension

Last updated: April 2007

Batch No: _____

Disclaimers: CAUTION: The information on The Hospital for Sick Children World Wide Web Site has been developed specifically for The Hospital for Sick Children. It is provided for informational purposes for qualified health care professionals. This material is not intended as a substitute for consulting qualified health care professionals. Patient circumstances will vary and some information may have become outdated as a result of more recent medical developments.

Ingredients	Mfr	Lot #	Expiry Date	Quantity	Measured	Checked
Sotalol HCl 160mg tablets	Bristol/ NOP			1		
Ora-Plus®	Paddock			16mL		
Ora-Sweet®	Paddock			16mL		
Ora-Plus®/Ora-Sweet®	SickKids			qs 32mL		

Equipment

- small mortar and pestle
- glass stirring rod
- 3 mL oral syringe x 1
- 30 mL graduated measure
- 50 mL amber bottle
- shrink seal x 1


✓ Procedure

1. Measure out Ora-Plus® and Ora-Sweet® and mix well.
2. Crush Sotalol tablets in the mortar with pestle to a fine powder.
3. Levigate the powder with small amounts of vehicle to form a paste, then add more vehicle until a liquid is formed. Transfer the contents into the graduate. Use additional vehicle to rinse the remaining drug from the mortar.
4. Make up to the final volume using the graduate and a 3mL syringe. Stir well.
5. Transfer the suspension into the amber bottle.
6. Label and affix shrink seal.

Storage: Room Temperature

Expiry: 60 days

Sample Label:

	SOTALOL HCL 5MG/ML ORAL SUSPENSION
L#:	EXP:
ROOM TEMPERATURE	SHAKE WELL (32mL)

Date and Signature (Technician and Pharmacist Final Check): _____
