

SickKids levoFLOXacin 50 mg/mL Oral Suspension

Batch No:

Ingredients	Mfr	Lot #	Expiry Date	Quantity	Measured	Checked
levofloxacin				10		
500 mg tablets						
Oral-Mix	Medisca			q.s.100 mL		

Additional Information:

Actavis/Teva or JAMP brands.

Equipment:

mortar and pestle	glass stirring rod	
graduated measure		

Procedure:

Follow your Dept. procedures for risk assessment/training/PPE/equipment/facilities/NAPRA level

- 1. Soak tablets in a small amount of vehicle for a minimum of 30 minutes.
- 2. Then levigate tablets by hand with a pestle for a minimum of 10 minutes to form a smooth paste. Continue to levigate as vehicle is added in small amounts until a liquid is formed.
- 3. Transfer liquid contents from mortar to graduate.
- 4. Use a small amount of vehicle to rinse mortar and add it to graduate.
- 5. Use vehicle to q.s. to the final volume. Stir well.
- 6. Transfer to amber bottle and label.

Quality Control:

Expected Product Appearance	Additional Notes	
White suspension	Original stability study was performed with SANDOZ brand	
	500mg tablets	

Storage:	Room temperature	
Packaging:	Amber glass/plastic PET bottles	
BUD:	90 days	

Sample Label:

	levoFLOXacin 50 mg/mL Oral Suspension			
SickKids Pharmacy	Lot:	BUD:		
	Room Temperature	Shake Well		

Date Made/Prepared By/Checked By: _____

Reference:

1. Tadrif V et al. (2020) Stability assessment of levofloxacin in three different suspension vehicles. Journal of Pharmacy Practice and Research. 50. 10.1002/jppr.1620.

Formulation Reviewed: December 2021

Disclaimer: The information in this compounding and preparation worksheet was developed for in-house use only by The Hospital for Sick Children ("SickKids") and its staff. While SickKids endeavours to keep this information up to date and accurate, patient circumstances will vary and some information may become outdated as a result of recent medical developments following the time of printing. Any reliance you place on such information is therefore strictly at your own risk. Please note that the doses provided do not necessarily conform to the manufacturer's recommendations, and formulary drugs may not be licensed for use in children. The reader is cautioned to check the product information sheet provided by the manufacturer, and to verify indications, contraindications and recommended dosages. SickKids make no representations or warranties of any kind, express or implied, about the completeness, accuracy, reliability, or suitability of the information and related products contained in this worksheet for any purpose. SickKids accepts no responsibility for the reader's use, misuse or misinterpretation of the information, and does not promise success in achieving any desired result. In no event will SickKids be liable for any damage, liability or loss or for any indirect, incidental or consequential damages arising from the use of this information.