



# IMPROVING THE VACCINATION EXPERIENCE:


## Using numbing cream to reduce pain in babies under one year of age

Numbing creams contain medicines called topical anesthetics. They are used to temporarily numb the skin and can be used to reduce the pain of vaccination.

You can buy numbing cream at your local pharmacy without a prescription. Talk to your pharmacist if you have any questions.


### WHERE TO APPLY NUMBING CREAM

Vaccinations are usually given in the upper leg (thigh). Before you apply the cream, make sure you confirm the correct area(s) for vaccination with a health-care provider. For [children one year of age and older](#) vaccinations are given in a different area of the body.



### HOW TO APPLY NUMBING CREAM

1. Check the label and expiration date of the cream before using it.
2. Decide which leg to apply the cream to. If your baby is having more than one vaccination on the same day, then you may have to apply the cream to both legs.
3. Do not apply the cream on an open wound such as a cut or scratch.
4. If opening a new tube of cream, use the point on the top of the cap to break the seal over the tube.
5. Locate the area on the upper outer thigh (as shown in the image) where you will apply the cream.
6. Apply between a nickel and a quarter size amount of cream to the skin (do not touch the cream or rub it in).
7. Cover the site by wrapping the upper leg with plastic wrap or use a clear bandage (such as Tegaderm®) to keep the cream in place.
8. If cream oozes out from under the covering, wipe it away with a tissue. Avoid contact with the mouth, eyes or nose. Rinse your hands with water after applying the cream.



Financial contribution from



Public Health  
Agency of Canada

Agence de la santé  
publique du Canada



Scan for more  
resources >



9. Mark the time of application. Make sure the cream is kept in place for the recommended wait time (30 to 60 minutes). See the table below for the wait times for some commercially available creams.
10. Just before vaccination, remove the covering and wipe away the cream with a tissue. Make sure the vaccination is given in the pre-treated area.

<b>Numbing cream</b>	<b>Brand names</b>	<b>Minimum time to take effect</b>
Lidocaine	Maxilene™ 4% or 5% Zensa™ 5%	30 minutes
Tetracaine	Ametop™ 4%	45 minutes
Lidocaine-prilocaine	EMLA™ 5% (also comes as a patch)	60 minutes

## **DOES NUMBING CREAM HAVE ANY SIDE EFFECTS?**

When used as directed, numbing cream has few side effects.

- Your baby’s skin may be whiter or redder for a short time. This is normal.

Call your health-care provider if:

- You get the cream in your or your baby’s mouth or eyes.
- If your baby shows signs of an allergic reaction to the cream. This includes rash, hives, swelling, wheezing or shortness of breath.

## **STORING NUMBING CREAM**

Store numbing cream in its original container out of children’s reach and away from direct sunlight, heat and humid places such as the bathroom. Make sure the cap is closed tightly and, if possible, lock the cream away with other medicines.